



OPEN ACCESS

Edited and reviewed by:

Mickaël Desvaux, Institut National de la Recherche Agronomique (INRA), France

*Correspondence: Shun Adachi f.peregrinusns@mbox.kyoto-inet.or.jp

[†]Present address:

Shun Adachi, Department of Microbiology, Kansai Medical University, Hirakata, Japan Yasuhiro Murakawa, RIKEN-IFOM Joint Laboratory for Cancer Genomics, RIKEN Center for Integrative Medical Sciences, Yokohama, Japan

[‡]Deceased

Specialty section:

This article was submitted to Microbial Physiology and Metabolism, a section of the journal Frontiers in Microbiology

> Received: 27 November 2020 Accepted: 11 December 2020 Published: 13 January 2021

Citation:

Adachi S, Murakawa Y and Hiraga S (2021) Corrigendum: Dynamic nature of SecA and Its associated proteins in Escherichia coli. Front. Microbiol. 11:634250. doi: 10.3389/fmicb.2020.634250

Corrigendum: Dynamic nature of SecA and Its associated proteins in *Escherichia coli*

Shun Adachi*[†], Yasuhiro Murakawa[†] and Sota Hiraga[‡]

Department of Radiation Genetics, Graduate School of Medicine, Kyoto University, Kyoto, Japan

Keywords: chromosome partition, SecA, SecY, AcpP, MukB, DNA topoisomerase

A Corrigendum on

Dynamic nature of SecA and Its associated proteins in Escherichia coli

by Adachi, S., Murakawa, Y., and Hiraga, S. (2015). *Front. Microbiol.* 6:75. *doi:* 10.3389/*fmicb.*2015.00075

In the original article, there were errors in **Figures 1C,F** and **G** as published. Images in the 1st and 2nd rows of Panel C and an image of DAPI in the 3rd row of Panel F are from different optical sections. For Panel G, the left column is from Native SecA, not Native SacA. The corrected **Figure 1** appears below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Copyright © 2021 Adachi, Murakawa and Hiraga. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.



FIGURE 1 Three-dimensional deconvoluted images of various immunostained proteins. Cells were grown in medium L at 30°C and fixed with 70% methanol. Red, immunofluorescence of GFP or native proteins. Green, DAPI staining. Merge, merged images of red and green images. Note that synthesized yellow signal is only recognizable with eyes when the ratio of each signal is between ~4:6 and 6:4. More difference is recognizable as mere green *(Continued)*

FIGURE 1 | or red. (**A**) Images of a DAPI-stained wild-type cell before and after 3D deconvolution in the Z-axis. (**B**–**F**) Fixed cells expressing GFP_{uv4}-fused proteins or GFP_{uv4} protein (not fused with any protein) were immunostained using an anti-GFP monoclonal antibody. Fluorescence images of the proteins were analyzed with 3D deconvolution at 0.5-μm intervals in the Z-axis. Images of DAPI-stained nucleoids were also analyzed with 3D deconvolution. The yellow arrowheads indicate the clustered localization of protein molecules. (**B**) SecA-GFP_{uv4}, (**C**) SecY-GFP_{uv4}, (**D**) AcpP-GFP_{uv4}, (**E**) GFP_{uv4} (not fused with any protein), (**F**) No GFP_{uv4}. (**G**–**I**) Deconvoluted images of DAPI-stained nucleoids were also deconvoluted images of DAPI-stained sing rabbit antisera for native proteins. Images of DAPI-stained nucleoids were also deconvoluted images of DAPI-stained nucleoids were also deconvoluted. (**G**) Native SecA, (**H**) Native SecY, (**I**) Native AcpP, (**J**) Three-dimensional deconvolution images of immunostained using rabbit antisera for native proteins.